

POPULATION CEFAZOLIN PHARMACOKINETICS

BEFORE, DURING AND AFTER CARDIOPULMONARY BYPASS IN CHILDREN UNDERGOING CARDIAC SURGERY

Pieter A.J.G. De Cock<sup>1,2</sup>, Hussain Mulla<sup>3</sup>, Karen Jacobs<sup>2</sup>, Kim Vanderburgh<sup>t2</sup>, Sarah Desmet<sup>1,2</sup>, Brett McWhinney<sup>4</sup>, Jacobus Ungerer<sup>4</sup>, Filip De Somer<sup>5</sup>, Annelies Moerman<sup>6</sup>, Hugo Robays<sup>1</sup>, Katrien Francois<sup>5</sup>, Peter De Paepe<sup>2</sup>

<sup>1</sup>Department of Pharmacy, Ghent University Hospital, Ghent , Belgium, <sup>2</sup>Heymans Institute of Pharmacology, Ghent, Belgium, <sup>3</sup>Department of Pharmacy, University Hospitals of Leicester, Leicester, UK, <sup>4</sup>Department of Chemical Pathology, Pathology Queensland, Brisbane, Australia, <sup>5</sup>Department of Cardiac Surgery, Ghent University Hospital, Ghent, Belgium, <sup>6</sup>Department of Anaesthesiology, Ghent University Hospital, Ghent, Belgium

Background

Scarce data are available to guide cefazolin dosing in children undergoing cardiac surgery with cardiopulmonary bypass (CPB).

Objectives

The objective of this trial is to derive a model-based dosing regimen for cefazolin in this patient population.

Methods

- 56 infants and children were included and received following intravenous dosing regimen: 25 mg/kg as a bolus 30 minutes before surgical incision, just before cessation of CPB, 8 hours after the 2nd dose and 8 hours after the 3rd dose.
- Blood samples were collected before, during and after cardiopulmonary bypass. Total (n=497) and unbound cefazolin concentrations (n=494) were measured using a validated High-Pressure Liquid Chromatography method; unbound concentrations were separated using a validated ultrafiltration method.
- NONMEM® version 7.2 was used for population PK modeling and covariate analysis: one, two and three compartmental models with first-order elimination were evaluated. R®, PsN® and Xpose® tools were used for data manipulation, graphical and statistical summaries.
- Glomerular Filtration Rates (eGFR) pre-, on- and post-CPB were estimated using the Modified Schwartz formula.
- Internal validation of the final model was performed using a non-parametric bootstrap (n=500 samples), a prediction corrected Visual Predictive Check (pcVPC) (n=1000 simulations) and the Normalised Prediction Distribution Error method (NPDE)(n=1000 simulations).
- To evaluate the current dosing regimen, Monte Carlo simulations (n=1000 simulations) were performed for a typical patient, procedure duration and median eGFR during the pre-, on- and post-CPB period. A minimum time frame of 60% of the dosing interval in which the unbound drug concentration remained above the Minimal Inhibitory Concentration (MIC) of *Staphylococcus aureus* (8 mg/L) was aimed for.

Intermediate results

PATIENT CHARACTERISTIC	MEDIAN (RANGE)
Age (days)	270 (5-5439)
Weight (kg)	6.8 (2.7-70)
Procedure duration (h)	2.15 (1.21-8.50)
CPB duration (h)	1.30 (0.37-5.1)
eGFR (mL/min/1.73 m2) pre CPB	108 (32-187)
eGFR (mL/min/1.73 m2) during CPB	80 (29-112)
eGFR (mL/min/1.73 m2) post CPB	74 (27-135)
Cardiac output (L/h) during CPB	44.82 (18.36-249)
Albumin (g/dL)	36 (21.8 -53)
Priming volume (mL)	175 (150-1000)

Table 1. Patient characteristics

Parameter	Estimate	RSE (%)	Median Estimate from Bootstrap (n=500 <sup>1</sup> )	2.5 <sup>th</sup> Percentile from Bootstrap (n=500 <sup>1</sup> )	97.5 <sup>th</sup> Percentile from Bootstrap (n=500 <sup>1</sup> )
Fixed effects					
$CL_{unbound} = CL_{pop} \times \left(\frac{WT}{WT_{med}}\right)^{0.75} \times \left(\frac{eGFR}{6.48}\right)$					
CL (L/h)	1.8	5	1.81	1.64	1.97
$Q_{unbound} = Q_{pop} \times \left(\frac{WT}{WT_{median}}\right)^{0.75}$					
Q1 (L/h)	4.79	6	4.84	4.24	5.45
Q2 (L/h)	FIXED to cardiac output during CPB				
$V_{unbound} = V_{pop} \times \left(\frac{WT}{WT_{median}}\right)^1$					
V1 (L)	1.87	6	1.88	1.63	2.10
V2 (L)	2.29	7	2.32	2.02	2.64
V3 (L)	FIXED to CPB priming solution volume				
$B_{max} = (\theta_{intercept} + \theta_{slope} \times \text{albumin concentration})$					
$\theta_{intercept}$ (mg/L)	132	13	133	98	169
$\theta_{slope}$	1.60	21	1.64	1.11	2.20
K <sub>D</sub> (mg/L)	47.7	10	48.5	40.3	59.2
Interindividual variability					
$\omega^2$ CL	0.095	28	0.088	0.046	0.15
$\omega^2$ V1	0.111	38	0.107	0.039	0.188
$\omega^2$ V2	0.241	25	0.232	0.116	0.365
$\omega^2$ Bmax	0.043	38	0.040	0.016	0.074
Residual variability (proportional)					
$\sigma^2$ (unbound concentration)	0.074	11	0.073	0.056	0.091
$\sigma^2$ (total concentration)	0.026	16	0.025	0.017	0.034

Table 2. Parameter estimates (<sup>1</sup> 273 successful runs)




Figure 2 : Goodness-of-fit plots and distribution histogram of NPDE for unbound concentrations




Figure 3: Goodness-of-fit plots and distribution histogram of NPDE for total concentrations




Figure 1 : Structure of the final model




Figure 4. Prediction-corrected Visual Predictive Check (n=1000) (left: panel unbound; right panel: total concentrations)

Legend: dots: observations; solid red line: observed median; dashed red lines: observed 5<sup>th</sup> and 95<sup>th</sup> percentiles; shaded areas : 95% CI of simulated 5<sup>th</sup>, 50<sup>th</sup> and 95<sup>th</sup> percentiles




Figure 5. Monte Carlo simulations for a typical patient, procedure duration and median eGFR pre-,on- and post CPB

Legend : solid line: median ; dotted lines : 5<sup>th</sup> and 95<sup>th</sup> percentiles (n= 1000); red dashed line: MIC cut-off

Discussion and Perspectives

- The proposed model adequately describes bound and unbound plasma cefazolin pharmacokinetics in infants and children undergoing cardiac surgery with cardiopulmonary bypass.
- Based on Monte Carlo simulations, the current dosing regimen needs to be optimised with a maximum dosing interval of 5-6h in postoperative patients.
- In a next step, a tissue concentration pharmacokinetic model will be developed to investigate the adequacy of dosing in terms of tissue penetration.